

Colonoscopy with SuPrep

Your Procedure:

You are scheduled for a colonoscopy, a procedure in which a doctor examines the lining of your large intestine by looking through a flexible tube called a colonoscope. If growths or other abnormalities are found during the procedure, the doctor may remove the abnormal tissue for closer examination or biopsy.

Procedure Date: _____

Arrival Time: _____ **Procedure Time:** _____

YOUR BOWEL PREP IS EXTREMELY IMPORTANT! PLEASE READ INSTRUCTIONS CAREFULLY.

1 Week Prior	3 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
<p>▶ Purchase the following: -SuPrep Bowel Prep Kit (prescription from Dr.) -Clear liquids (no red or purple – see list for clear liquid choices) ▶ Optional – for sore bottom: -Plain or Aloe Baby Wipes -Desitin or A&D ointment ▶ Arrange for a responsible adult to stay with you during the procedure and take you home post-procedure. ▶ If you take fiber supplements or medications containing iron discontinue those 7 days before your appointment. This includes multi-vitamins with iron. ▶ You may continue your “baby” aspirin and regular aspirin. Let us know if you are on Plavix, coumadin, or other blood thinners. ▶ If you have diabetes you have been given a Diabetic Prep Sheet. ▶ Stop taking these medications:</p> <p>_____</p> <p>_____</p>	<p>▶ Begin low fiber diet. See handout for details.</p> <p>▶ Stop consuming all high-fiber foods.</p> <p>▶ Last chance to cancel appointment. Call your physician’s office at the number below if you need to reschedule your procedure or if you have any questions:</p> <p>_____</p>	<p>▶ Continue low fiber diet.</p> <p>▶ Drink at least 4 large glasses of water throughout the day.</p> <div data-bbox="884 862 1052 915" style="text-align: center;">  </div> <p style="text-align: center;">No solid foods after midnight!</p>	<p>▶ Begin clear liquid diet.</p> <p style="text-align: center;">See clear liquid choice handout for suggestions and recipes!</p> <p>▶ Drink at least 4 large glasses of water throughout the day.</p> <div data-bbox="1234 862 1402 915" style="text-align: center;">  </div> <p>▶ 5pm: Pour 1 six (6) oz bottle of SuPrep liquid into the mixing container. Add cold water to the 16 oz line on the container. Drink ALL of the liquid.</p> <p>▶ 6-7pm: Drink 2 more 16 oz containers of cool/cold water over the next hour.</p> <p>If you experience nausea or vomiting take a 15-30 minute break, then continue drinking prep solution.</p>	<p style="text-align: center;">2nd Part of Split Prep</p> <p>▶ 5-6 hours before your procedure: Pour 1 six (6) oz bottle of SuPrep liquid into the mixing container. Add cold water to the 16 oz line. Drink ALL of the liquid.</p> <p>▶ Drink 2 more 16 oz containers of cool/cold water over the next hour. (NOTE: This should be completed at least 4 hours prior to your procedure.)</p> <p>▶ After your prep, the results should appear <u>clear yellow</u> or <u>clear green liquid</u> with <u>no solid matter</u>. If not, please call your doctor at the office or medical exchange.</p> <p>▶ NOTHING by mouth at least 4 hours prior to your scheduled procedure time – including gum, tobacco, hard candy/mints!</p> <p>▶ You may take your essential morning medications with a few small sips of water, at least 4 hours prior to your procedure, unless otherwise directed by your physician.</p> <p>▶ Be sure to bring the following:</p> <ul style="list-style-type: none"> -Responsible adult driver to stay with you and drive you home. Your procedure will not start without your driver present. -Insurance Cards -Driver’s License/Photo ID -Any co-insurance fees due